



Pectus Carinatum Repair by Minimally Invasive Procedure New Trends About Pectus Carinatum Repair

New Trends Pectus Repair

Horacio Abramson¹, Sebastián Pisoni², Manuel Rodríguez², Gustavo Segura²

¹ Htal. Antonio Cetrángolo, Vte. López, Buenos Aires, Head of Thoracic Surgical Service, ²Thoracic Surgical Service, Htal. Antonio Cetrángolo, Argentina.

Introduction

The aim of this article is to show a new minimally invasive procedure for Pectus Carinatum Repair. Moreover the purpose is also to mention the re-

Background

Since the 1988 published article that introduced the Nuss Procedure for Pectus Excavatum correction [6], interest has emerged among surgeons dedicated to chest malformation diseases to perform the Pectus Carinatum Repair with less invasive methods.

This trend has been increased starting with the original work we showed in 2005 [7]. Through a novel technique and using an original prosthesis (Patent no. US 7,156,847 B2), this procedure corrects the pectus carinatum (p c) by means of the insertion of a compressive pre-sternal implant.

Later the results of the first 40 corrected patients were shown at the American Pediatric Surgical Association (APSA) Congress, in Phoenix, AZ USA, 2007. [8]

The original procedure has been described by Steven Rothemberg M.D.[9] as a "Reverse Nuss". We can also

cent contributions developed by experts in this realm to reduce the aggressiveness of the open and invasive techniques utilized during the last half century [1-5].

schematically depict it as an "Internal Orthosis".

The novelty takes advantage of the therapeutic experience of pre-sternal chest braces [10, 11], which correct the anterior chest wall protrusions without surgery: avoiding costo-chondral resections, sternal osteotomies, wide muscle and dermical flaps, and wide skin incisions that can provide frequent keloids. On the other hand this innovation avoids the lack of adherence to treatment which is an important factor for orthotic failure treatments.

It is known that the final favorable brace appliance result lies in systematic, continuous and progressive increases of leaning on the back and front chest.

Many factors create practical difficulties for long term application: the annoyance in having to use it many hours a day; troubles in warm weather countries and dermical damage over the protruded zones among other findings.

DOI: 10.4328/JCAM.10.1.41 Received: 17.11.2009 Accepted: 26.11.2009 Printed: 01.01.2010 J.Clin.Anal.Med. 2010;1.1:60 - 3

Corresponding author: Dr. Horacio Abramson. Av. Libertador 1688, Vicente López, Pcia. de Buenos Aires. CP: 1638 Buenos Aires, ARGENTINA.

E-mail: habramson@intramed.net.ar

The New Apparatus and Procedure

This innovation is comprised of a new technique and a new apparatus for the correction of Pectus Carinatum by minimally invasive procedure. In this technique the sternochondral region is compressed by implanting a metal bar in the pre-sternal region and securing it bilaterally to the posterolateral portion of the costal arches (Figure 1).

The target patients are children and youths suffering the Pectus Carinatum Syndrome both in the symmetric or asymmetric forms (Figure 2, 3). Adequate elasticity of the chest is a key factor required in this surgical procedure to achieve an effective molding of the chest wall. The ideal timeframe for the operation is the

Discussion and Conclusions

Pectus Carinatum is a frequent malformation and creates many medical consults because it is difficult to hide and deeply affects personality and behaviour. The orthotic procedures are cumbersome and time-consuming and generally have poor results. The surgical procedures have shown good results since the Ravitch technique. Open techniques are aggressive and leave scars that often bring keloids. This new method has the following advantages: no extensive incisions, no need for long muscle flaps, no resection of chondral or costal arches; no sternal section. This technique does not affect breathing mechanisms by avoiding the post-surgical rigidity created by extensive resections in the thoracic wall. This method brings about very good thoracic correction results. It is a new procedure similar to Nuss' method for Pectus Excavatum repair but the reverse. Moreover, the apparatus is useful to correct Pectus Excavatum.

The aligned holes at both tips of the new designed bar

Novelties about Pectus Carinatum Techniques (Less Invasive Procedures)

In 1997 [12] Dr. Kobayashi from Japan reported a technique by endoscopy assistance, but including cartilage resections and sternal ostectomy by means of a little pre-sternal incision.

In 2004 [13] Dr. Fonkalsrud Eric proposed a technique with minimally cartilage resections that keeps characteristics of open surgeries: wide incisions, muscle flaps, osteo-chondral resections and sternal fracture.

In 2005 [14, 15] Dr. Saxena reports a less invasive pro-

cedure for both excavatum and carinatum disease: the Willital technique also known as PLIER, (Pectus Less Invasive Extra pleural Repair).

period of rapid body growth. The compression system acts in a way similar to that of orthodontic braces. The results considered in terms of the contour of the chest wall and their anatomical characteristic is comparable to invasive techniques. The late development of chest wall rigidity that may occur when parietal resection is performed is also avoided.

Two fixing plates are firmly implanted anchored bilaterally by pericostal stainless steel wires sutures. The link between the fixation plates and the compressive bar is with screws. The system remains implanted until ossification is achieved, and an adequate thoracic shape is obtained (Figure 4, 5). The bar is removed at a later time.

let the surgeon apply progressive pressure at both sides finding the best possible thoracic shape. The threaded holes along the groove of the fixation plate make this useful manoeuvre easy and safe.

This technique has been inspired by the Nuss Procedure. However, this procedure has important differences. The extra thoracic, pre-sternal position of the implant decreases the risks of serious operative accidents and makes thoracoscopy unnecessary. Early discharge, rapid return to activity and freedom from positional restrictions, no need for long periods of avoiding rotation or displacement of the implant are the principal achievements of the innovation.

Asymmetric conditions are not a technical barrier with this minimally invasive technique. The pressure on the sternum and the protruded region of the chondrosternal junctions is transferred through the costal arches to a fixed point (the vertebral backbone); thus, the asymmetries are corrected. Tolerance of the implant was good especially in young patients younger than 14 years.

cedure for both excavatum and carinatum disease: the Willital technique also known as PLIER, (Pectus Less Invasive Extra pleural Repair).

However, the PLIER technique maintains the characteristics of the open techniques: chondro-costal resections, sternal ostectomies, etc.

In 2006 [16] Dr. Klaus Schaarschmidt published another less invasive technique similar to the Kobayashi presentation, under assisted endoscopy, but including chondral resections and sternotomies.

The Hungarian experience developed by Dr. Attila Kal-



Figure 1. Squeme of the implant

man [17] and Andras Hock [18] after they were aware of our technique in 2005 which was published in the *Separ Bronconeumology Review*. Both reports are similar. The bar is implanted in intra thoracic position, and then pre-sternally and finally inserted intra-tho-



Figure 2. Patient before correction.

racily in the opposite hemithorax.

In spite of the favorable results, in my opinion this procedure has the disadvantage of being invasive in both pleural cavities. Moreover, the implant stability depends on the wavy shape that the compressor bar adopts. For that reason, the removal of the bar may be more difficult or risky due to this particular design.

In 2008 [19] Dr. Seock Yeol Lee from Korea, reported his experience performing partial cartilage resection and pre-sternal compression with a stainless steel bar




Figure 3. XRay before correction.

in 10 patients with pectus carinatum.

In 2009 [20] Dr. Sunghoon Kim reported his experience from Children's Hospital Oakland, CA, USA. This technique was applied just in an unilateral Pectus Carinatum patient and has the disadvantage of being invasive making the cartilage resections within the thoracic cavity.

The Turkey experience was lead by Dr. Mustafa Yüksel from Istanbul [21]. He was the first surgeon to give importance to the new procedure which he found out about at the European Association for Cardio-Thoracic Surgery (EACTS) Congress in Barcelona 2005, and proceeded to apply it. Later in Buenos Aires in 2007 attending the World Federation of Associations of Pediatric Surgeons (WOFAPS) Congress showed his first experience. After 3



Figure 4. Patient after correction.



Figure 5. XRay after correction.

Pectus Carinatum repairs decided to change the Nuss Bar for a design similar to the original implant.

In 2008 [22] Dr. Patricio Varela from Chile reported two cases corrected also by means of the compressive technique but utilizing the Nuss Pectus Excavatum design. The follow up was short: 9 and 3 months respectively, but showed the disengagement of the implant in one of the two patients.

References

- 1 Ravitch MM. The operative unusual sternal deformity with cardiac Symptoms. Operative correction. *J Thorac Surg* 1952;23:138-44.
- 2 Ravitch MM. The operative correction of pectus carinatum (pigeon breast). *Ann Surg* 1960;151:705-14.
- 3 Lester CW. Pigeon breast (pectus carinatum) and other protrusion deformities of the chest of developmental origin. *Ann Surg* 1953;137: 482-9.
- 4 Chin EF. Surgery of the funnel chest and congenital sternal prominence. *Br J Surg* 1957;186:360-76.
- 5 Brodtkin HA. Pigeon breast—congenital chondrosternal prominence. Etiology and surgical treatment by xyphosternopexy. *Arch Surg* 1958; 77:261-70.
- 6 Nuss D, Kelly RE, Croitoru DP, Katz ME. A 10-year review of a minimally invasive technique for the correction of pectus excavatum. *J Ped Surg.* 1988;33:545-52.
- 7 Abramson H. A minimally invasive technique to repair pectus carinatum. Preliminary report. *Arch Bronconeumol* 2005;41:349-51.
- 8 Abramson H, D'Agostino J, Wuscovi S. A 5-year experience with a minimally invasive technique for pectus carinatum repair. *J Ped Surg.* 2009; 44:118-124.
- 9 Rothenberg S. Current concepts in surgical management of pediatric lung and chest wall diseases. *Curr Opin Pediatr.* 2006;18:282-286.
- 10 Haje SA, Bowen JR. Preliminary results of orthotic treatment of pectus deformities in children and adolescents. *J Pediatr Orthop* 1992;12:795-800.
- 11 Vidal J, Nakach G. Tratamiento ortopédico de las deformaciones torácicas. In: Villadot R, Cohí O, Clavell S, editors. *Ortesis y prótesis del aparato locomotor. Columna vertebral.* Masson: Barcelona; 1994.85-93.
- 12 Kobayashi S, Yoza S, Komuro Y, Sakai Y, Ohmori K. Correction of pectus excavatum and pectus carinatum assisted by the endoscope *Plast Reconstr Surg.* 1997;99:1037-45
- 13 Fonkalsrud E, Anselmo D, Less Extensive Techniques for Repair of Pectus Carinatum. *The Undertreated Chest Deformity J Am. Coll. Surg.* 2004;198, (6) 898-905.
- 14 Saxena AK. Pectus excavatum, pectus carinatum and other forms of thoracic deformities. *J. Ind. Ass. Ped. Sur.* 2005; Vol.10, No.3:147-57.
- 15 Saxena AK. Pectus less invasive extrapleural repair (PLIER). *J Plast Reconstr Aesthet Surg.* 2009;62,5:(6)63-8.
- 16 Schaarschmidt K, Kolberg-Schwerdt A, Lempe M, Schlesinger F. New Endoscopic Minimal Access Pectus Carinatum. *Ann Thorac Surg* 2006;81:1099 -104.
- 17 Kálmán A. Initial results with minimally invasive repair of pectus carinatum., *Thorac Cardiovasc Surg.* 2009.
- 18 Hock A, Minimal access treatment of pectus carinatum: a preliminary report. *Pediatr Surg Int.*; 2009. (article in press).
- 19 Lee SY, Oh JY, Lee SJ, Lee CS. A Modified Technique for Pectus Carinatum Surgery: Partial Costal Cartilage Resection and Pre-sternal Compression with Using a Stainless Steel Bar *Korean J Thorac Cardiovasc Surg* 2008;41:742-6.
- 20 Sunghoon Kim, Olajire Idowu. Minimally invasive thoracoscopic repair of unilateral pectus carinatum. *J. Ped. Sur.* 2009; 44, 471-4.
- 21 Yuksel M, Onen A, Bostanci K, Sanli A, Lacin T, Akgul A G, Batirel H F. Early results following the minimally invasive repair of pectus carinatum deformity using a newly designed bar: a two institution experience. *17th European Conference on General Thoracic Surgery* 31 May-3 June 2009, Krakow, Poland.
- 22 Varela P, Leopold E. Corrección mínimamente invasiva de pectus carinatum con prótesis compresiva (operación de Abramson). *Revista Pediatría Electrónica* 2008,(5)2.